

REVIEW

Outcome of primary root canal treatment: systematic review of the literature – Part 1. Effects of study characteristics on probability of success

Y.-L. Ng¹, V. Mann², S. Rahbaran¹, J. Lewsey³ & K. Gulabivala¹

¹Unit of Endodontology, UCL Eastman Dental Institute, University College London; ²Department of Medical Statistics, London School of Tropical Medicine and Hygiene; and ³Clinical effectiveness Unit, The Royal College of Surgeons of England, London, UK

Abstract

Ng Y-L, Mann V, Rahbaran S, Lewsey J, Gulabivala K. Outcome of primary root canal treatment: systematic review of the literature – Part 1. Effects of study characteristics on probability of success. *International Endodontic Journal*, 40, 921–939, 2007.

Aims The aims of this study were (i) to conduct a comprehensive systematic review of the literature on the outcome of primary (initial or first time) root canal treatment; (ii) to investigate the influence of some study characteristics on the estimated pooled success rates.

Methodology Longitudinal clinical studies investigating outcome of primary root canal treatment, published up to the end of 2002, were identified electronically (MEDLINE and Cochrane database 1966–2002 December, week 4). Four journals (*International Endodontic Journal*, *Journal of Endodontics*, *Oral Surgery Oral Medicine Oral Pathology Endodontics* and *Dental Traumatology & Endodontics*), bibliographies of all relevant papers and review articles were hand-searched. Three reviewers (Y-LN, SR and KG) independently assessed, selected the studies based on specified inclusion criteria, and extracted the data onto a pre-designed proforma. The study inclusion criteria were: longitudinal clinical studies investigating root canal treatment outcome; only primary root canal treatment carried out on the teeth studied; sample size

given; at least 6-month postoperative review; success based on clinical and/or radiographic criteria (strict, absence of apical radiolucency; loose, reduction in size of radiolucency); overall success rate given or could be calculated from the raw data. The findings by individual study were summarized and the pooled success rates by each potential influencing factor were calculated for this part of the study.

Results Of the 119 articles identified, 63 studies published from 1922 to 2002, fulfilling the inclusion criteria were selected for the review: six were randomized trials, seven were cohort studies and 48 were retrospective studies. The reported mean success rates ranged from 31% to 96% based on strict criteria or from 60% to 100% based on loose criteria, with substantial heterogeneity in the estimates of pooled success rates. Apart from the radiographic criteria of success, none of the other study characteristics could explain this heterogeneity. Twenty-four factors (patient and operative) had been investigated in various combinations in the studies reviewed. The influence of preoperative pulpal and periapical status of the teeth on treatment outcome were most frequently explored, but the influence of treatment technique was poorly investigated.

Conclusions The estimated weighted pooled success rates of treatments completed at least 1 year prior to review, ranged between 68% and 85% when strict criteria were used. The reported success rates had not improved over the last four (or five) decades. The quality of evidence for treatment factors affecting primary root canal treatment outcome is sub-optimal; there was substantial variation in the study–designs. It

Correspondence: Dr Y.-L. Ng, Unit of Endodontology, UCL Eastman Dental Institute, UCL, 256 Grays Inn Road, London WC1X 8LD, UK (Tel.: 020 7915 1233; fax: 020 7915 2371; e-mail: p.ng@eastman.ucl.ac.uk).

would be desirable to standardize aspects of study–design, data recording and presentation format of outcome data in the much needed future outcome studies.

Keywords: outcome, root canal treatment, success, systematic review.

Received 27 March 2007; accepted 6 July 2007

Introduction

There has been a surge of interest in formulating clinical guidelines for optimal treatment of diseases based on properly conceived and executed research. The gold standard for informing clinical practice is putatively the randomized controlled trial (RCT); however, neither medical nor dental practice has been generally well supported by such evidence. Sackett *et al.* (1996) now famous definition, ‘the conscientious, explicit and judicious use of current best evidence in making decisions about the care of individual patients’, not only embraces the notion of grades of evidence but also recognizes that optimal levels of evidence may not be available for all situations. There is therefore a need to synthesize an objective over-view based on *available* evidence. Depending upon the quality and quantity of the data, systematic reviews can be of several different kinds: traditional reviews; meta-analysis leading to an estimate of effect size; best evidence synthesis; and the hypothetico-deductive approach, in which the effort is directed at evaluating the evidence for and against a given theory, in an attempt to solve the problem of why contradictory results appear, rather than simply averaging often incompatible data (Eysenck 1994).

For the outcome of endodontic treatment, there are eight published systematic reviews, which have used different approaches in synthesis of information from the literature. Basmadjian-Charles *et al.* (2002) and Paik *et al.* (2004) used a systematic approach for literature search but a traditional approach for evaluating the variables impacting on the success and failure of the root canal retreatment. Two reviews (Hepworth & Friedman 1997, Peterson & Gutmann 2001) calculated the weighted-average success rates by each factor under investigation. Neiderman & Theodosopoulou (2003) estimated the number needed to treat when comparing two types of treatments. Three reviews (Lewsey *et al.* 2001, Kojima *et al.* 2004, Sathorn *et al.* 2005) estimated the size of effect of individual factors which included presence of preoperative pulpal and periapical status, apical extent of root filling & number of treatment visits, using meta-analysis. Except for Lewsey *et al.* (2001), none have investigated the

influence of study characteristics such as radiographic criteria for determination of treatment outcome and year of publication, on the data heterogeneity.

In the absence of sufficient gold standard level data, there is a need to synthesize ‘sub-standard’ data but there is a lack of formal guidelines to achieve this. In the absence of such guidelines, the authors proposed the use of a process of ‘triangulation’ of different analytical approaches as a sensible strategy. The purpose of this systematic review and synthesis was to: (i) identify the probable dominant factors influencing outcome; (ii) help prioritize the questions that need to be addressed; (iii) inform the design and data collection protocol for future RCTs. The outcome of this analysis will be presented in two parts.

The aim of the first part of this paper is to present the estimated pooled success rates of primary root canal treatment by aspects of study characteristics: decade of publication, study-specific criteria for success, unit of outcome measure, duration after treatment, geographical location of study and qualification of the operator.

Materials and methods

Literature search

Longitudinal clinical studies investigating the outcome of primary root canal treatment that were published up to the end of 2002 were identified electronically (MEDLINE database 1966–2002 December, week 4) using six keywords (root canal treatment, root canal therapy, endodontic treatment, endodontics, treatment outcome and success) and eight strategies (1 AND 5, 1 AND 6, 2 AND 5, 2 AND 6, 3 AND 5, 3 AND 6, 4 AND 5 and 4 AND 6). A Cochrane Library search was also conducted. PubMed was independently searched using the ‘related articles’ feature. Four journals (International Endodontic Journal, Journal of Endodontics, Oral Surgery Oral Medicine Oral Pathology Endodontics Radiology and Dental Traumatology & Endodontics) and bibliographies of all relevant papers and review articles were hand-searched. Unpublished studies were identified by

searching abstracts and conference proceedings. Personal contacts were also used to identify ongoing or unpublished studies. Full articles were obtained for all relevant titles identified through either electronic or other search methods.

Study selection, quality assessment and data extraction

Three reviewers (Y-LN, SR and KG) independently assessed and selected the studies based on the following inclusion criteria:

1. Clinical study on primary root canal treatment.
2. Stratified analysis of primary root canal treatment available, if root canal re-treatment cases had been included.
3. Sample size given.
4. At least 6-month postoperative review.
5. Success based on clinical and/or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria.
6. Overall success rate given or could be calculated from the raw data.
7. Presentations in English, German, Chinese and Japanese languages were accepted.

Disagreements on study inclusion were resolved by discussion. The reasons for study rejection at this or subsequent stages were recorded.

Data were extracted by all three reviewers independently using custom-designed data collection forms. The data collection form was piloted on several papers and modified for optimal utility before final use. The data extracted could be classified into six groups; success rates, study characteristics, demographic data of patients, pre-, intra- and postoperative factors. Any disagreement was discussed and data were excluded if agreement could not be reached.

Estimation of pooled success rates

STATA version 9.2 statistical software (StataCorp. College Station, TX, USA) was used to perform all statistical analyses. Un-weighted pooled success rate by each factor was calculated by dividing the total number of successful units with the total number of units within the respective category (according to Hepworth & Friedman 1997). In addition, the weighted pooled success rates were estimated using random effects meta-analysis with DerSimonian and Laird's methods (DerSimonian & Laird 1986). Statistical heterogeneity amongst the studies was assessed by Cochran's (Q) test (Cochran 1954).

Meta-regression models (Thompson & Higgins 2002) were used to explore the potential sources of statistical heterogeneity and to assess the effect of factors on estimating the pooled success rate. Factors related with study characteristics considered in the meta-regression analyses as covariates (and their sub-categories) were: decade of publication, study specific criteria (radiographic, combined radiographical & clinical) for success, unit of outcome measure (tooth and root), duration after treatment when assessing success ('at least 4 years' or '<4 years'), geographical location of the study (North American, Scandinavian and other countries), qualification of the operator (undergraduate students, postgraduate students, general dental practitioners (GDP), specialist or mixed group). If either the estimated proportion of total variation because of heterogeneity across studies (I^2) or the estimated between-study variance (τ^2) from the meta-regression model without covariate in the model was reduced substantially (>10%) when a covariate was included into the model, the respective covariate was considered to be a potential source of heterogeneity.

Results

A total of 119 papers were identified in the initial search, 51 articles were excluded for the reasons given in Table 1. Some papers presented different parts of the same study, so their data were combined for analyses in this review: (i) Heling & Tamshe (1970, 1971); (ii) Barbakow *et al.* (1980a,b, 1981); (iii) Morse *et al.* (1983a,b,c); (iv) Ørstavik *et al.* (1987) & Eriksen *et al.* (1988). Conversely, Kerekes (1978) presented two separate data sets in their paper, and were therefore considered as two separate studies in this review. As a result, 63 studies fulfilling the inclusion criteria were selected for this review. The year of publication of the selected studies ranged from 1922 to 2002 with the highest number of studies published in the 1980s ($n = 16$) (Table 2).

Each reviewer had entered 174 data points per selected study and the initial agreements amongst the three reviewers were moderate ($\kappa = 0.57-0.61$). As per protocol, following discussion about any disagreements, there was 100% concurrence on used data.

Methodological characteristics of included studies

Of the 63 studies included in this review, six were RCTs (Table 2). Others were cohort studies ($n = 8$) or retrospective observational studies ($n = 49$). Although

Table 1 Reasons for exclusion of the 51 articles

Article	Inclusion criteria (1–6) not fulfilled or other reasons for exclusion
Grove (1921)	¹ Clinical study for primary root canal treatment
Hinman (1921)	³ Sample size given ⁶ Overall success rate given or could be calculated
Grove (1923)	¹ Clinical study for primary root canal treatment
Coolidge (1926)	⁶ Overall success rate given or could be calculated
Rhein <i>et al.</i> (1926)	⁴ At least 6-month postoperative review
Puterbaugh (1926)	³ Sample size given ⁶ Overall success rate given or could be calculated
Hall (1928)	¹ Clinical study for primary root canal treatment
Appleton (1932)	⁴ At least 6-month postoperative review
Buchbinder (1936)	² Stratified analysis available
Macphee (1936)	⁶ Overall success rate given or could be calculated
Strindberg (1956)	² Stratified analysis of one RCT available
Frostell (1963)	² Stratified analysis of one RCT available
Nichols (1963)	¹ Clinical study for primary root canal treatment
Grossman <i>et al.</i> (1964)	² Stratified analysis of one RCT available
Engström <i>et al.</i> (1964)	² Stratified analysis of one RCT available
Ingle <i>et al.</i> (1965)	² Stratified analysis of one RCT available
Curson (1966)	¹ Clinical study for primary root canal treatment
Oliet & Sorin (1969)	⁵ Success based on clinical and/or radiographic criteria
Storms (1969)	² Stratified analysis of one RCT available
Ratliff (1973)	⁵ Success based on clinical and/or radiographic criteria
Cvek <i>et al.</i> (1976)	² Stratified analysis of one RCT available
Adenubi (1978)	Same data set as Adenubi & Rule (1976)
Taintor <i>et al.</i> (1978)	¹ Clinical study for primary root canal treatment
Vernieks & Messer (1978)	² Stratified analysis of one RCT available
Kerekes & Tronstad (1979)	² Stratified analysis of one RCT available
Markitziu & Heling (1981)	¹ Clinical study for primary root canal treatment
Hession (1981)	² Stratified analysis of one RCT available
Thoden van Velzen <i>et al.</i> (1981)	Same data set as Kerekes & Tronstad (1979)
Ashkenaz (1984)	⁵ Success based on clinical and/or radiographic criteria
Seto <i>et al.</i> (1985)	⁵ Success based on clinical and/or radiographic criteria
Ørstavik <i>et al.</i> (1986)	⁶ Overall success rate given or could be calculated
Teo <i>et al.</i> (1986)	² Stratified analysis of one RCT available
Kullendorff <i>et al.</i> (1988)	¹ Clinical study for primary root canal treatment
Molven & Halse (1988)	² Stratified analysis of one RCT available Same data set as Halse & Molven (1987)
Augsburger & Peters (1990)	⁴ At least 6-month postoperative review ⁵ Success based on clinical and/or radiographic criteria
Stabholz (1990)	¹ Clinical study for primary root canal treatment
Wong <i>et al.</i> (1992)	⁵ Success based on clinical and/or radiographic criteria
Ørstavik & Hörsted-Bindslev (1993)	⁶ Overall success rate given or could be calculated
Gutknecht <i>et al.</i> (1996)	⁴ At least 6-month postoperative review
Friedman (1997)	¹ Clinical study for primary root canal treatment
Ricucci & Langeland (1997)	¹ Clinical study for primary root canal treatment
Weine & Buchanan (1997)	¹ Clinical study for primary root canal treatment
Shi <i>et al.</i> (1997)	⁵ Success based on clinical and/or radiographic criteria
Weiger <i>et al.</i> (1998)	¹ Clinical study for primary root canal treatment
Caplan & White (2001)	⁵ Success based on clinical and/or radiographic criteria
Oliver & Abbott (2001)	¹ Clinical study for primary root canal treatment
Waltimo <i>et al.</i> (2001)	⁶ Overall success rate given or could be calculated
Lazarski <i>et al.</i> (2001)	⁵ Success based on clinical and/or radiographic criteria
Lynch <i>et al.</i> (2002)	⁵ Success based on clinical and/or radiographic criteria
Caplan <i>et al.</i> (2002)	⁵ Success based on clinical and/or radiographic criteria
Murakami <i>et al.</i> (2002)	⁴ At least 6-month postoperative review

nine studies (Grahnen & Hansen 1961, Storms 1969, Selden 1974, Heling & Kischinovsky 1979, Pekruhn 1986, Sjögren *et al.* 1990, Friedman *et al.* 1995, Chugal *et al.* 2001, Hoskinson *et al.* 2002) had included previously root-filled teeth in their sample, they had provided stratified analysis for primary treatment.

The recall rates (percentage of patients attending for follow-up after treatment) were reported by 39 studies and ranged from 11% to 100% with a median of 52.7%. Either root (27 studies) or tooth (36 studies) was used as the unit of outcome measure. The sample sizes ranged from 22 to 2921 teeth or 38 to 2921 roots; some studies only included single-rooted teeth, hence the number of teeth and roots were the same.

The treatment outcome was determined by radiographic examination alone (27 studies) or in combination with clinical findings (36 studies) (Table 2). Different radiographic criteria of success have been used and these were divided into: 'strict' (complete resolution of peri-apical lesion at recall) or 'loose' (reduction in size of existing peri-apical lesion at recall). For the radiographic assessment of the outcome of treatment, only 19 studies (Table 2) employed at least two observers to carry out the assessment. Observer(s) were calibrated prior to evaluation of radiographs in eight studies and intra- or inter-observer reliability tests were carried out in nine studies (Table 2).

Different studies have evaluated the influence of a range of different clinical prognostic factors on outcome but the combinations of factors reported vary (Table 3). The statistical methods used for analysing the association between potential influencing factors and treatment outcome were the chi-square test (31 studies), relative incidence distribution (two studies), logistical regression models (three studies), ANOVA (two studies), survival analysis (one study) and logistic regression models using generalized estimating equations (one study) (Table 2). Twenty-three studies did not analyse the data statistically or did not present such information.

Success rates by study characteristics

Outcome measure used

The reported success rates of root canal treatment ranged from 31% to 96% based on strict criteria and from 60% to 100% based on loose criteria. The weighted pooled success rates from studies using 'strict' criteria (data available from 40 studies) were about 10% lower than those from studies using 'loose' criteria (data available from 38 studies) regardless of examination method used (Table 4). Some studies ($n = 14$)

presented the success rates stratified by both strict and loose criteria.

After combining the data from the two examination methods, the pooled success rates estimated by meta-analyses were 74.7% (95% CI: 69.8–79.5%) from 40 studies using strict radiographic criteria and 85.2% (95% CI: 82.2–88.3%) from 36 studies using loose radiographic criteria. The estimated success rates by individual studies as well as the weighted pooled success rates by the two radiographic criteria are presented as Forrest plots in Figs 1 and 2. Meta-regression analyses showed the reported success rates based on strict radiographic criteria were 10.5% (4.4–16.7%, $P = 0.001$) lower than the success rates based on loose radiographic criteria. The radiographic criteria were also found to be responsible for part of the statistical heterogeneity, therefore the estimated success rates by individual factors were calculated separately for data based on the use of strict or loose criteria.

Duration after treatment completion

Most studies did not standardize the duration after treatment completion when the outcomes were reviewed, which ranged from 6 months to 30 years. Only 15 studies (Table 2) followed-up all the cases for at least 4 years. Attempts to pool data on success rates by different follow-up durations are confounded by the relatively small study numbers in some groups and may have produced distorted results. When strict criteria were used, the pooled success rates increased with longer follow-ups; the substantial increases were between 6 and 12 months and between 24 and 36 months after treatment (Table 4). However, there was no obvious trend in success rate by duration after treatment when loose criteria were used.

Year of publication

The pooled success rates based on 'loose' outcome criteria for each decade since the 1920s appeared to be similar with the highest pooled success rate at 88.2% during the 1980s (Table 4). However, the pooled success rates based on 'strict' outcome criteria for studies published during 1960s (79.7%) and 1970s (79.0%) were the highest. More importantly, the expected trend of progressively increasing success rates over the last century was clearly not in evidence.

Geographical location of study

About one-third of the studies were carried out in the USA or Canada (24 studies) and the rest were carried

Table 2 Study characteristics

Author (year)	Geographic location of study	^a Study-design	Recall rate (%)	^b Unit of measure	Sample size	^c Assessment of success	^d Radiographic criteria of success	^e Radiographic ≥2 observers	Calibration	Reliability test	^f Statistical analysis
Blayney (1922)	USA	R	28	T	104	C&R	L				
Auerbach (1938)	USA	R	22	T	211	C&R	L				
Buchbinder (1941)	USA	R	-	Ro	245	Ra	S				
Morse & Yates (1941)	USA	R	-	T	265	Ra	L				
Castagnola & Orlay (1952)	Scandinavia	R	68	T	1000	C&R	S	✓			
Grahnén & Hansen (1961)	Scandinavia	R	44	Ro	1277	C&R	S				
Seltzer et al. (1963)	USA	C	-	Ro	2921	Ra	L				
Zeldow & Ingle (1963)	USA	C	-	T	42	C&R	L				
Bender et al. (1964)	USA	R	30	Ro	706	Ra	L	✓			
Engström & Lundberg (1965)	Scandinavia	R	74	Ro	181	Ra	S				
Harty et al. (1970)	UK	R	60	Ro	1139	C&R	S				χ^2
Heling & Tamshe (1970, 1971)	Israel	R	27	T	213	C&R	S				
Cvek (1972)	Scandinavia	R	-	Ro	55	Ra	S				ANOVA
Selden (1974)	USA	R	11	T	556	Ra	L				χ^2
Werts (1975)	USA	R	23	T	47	C&R	S				χ^2
Adenubi & Rule (1976)	UK	R	-	Ro	870	C&R	S				χ^2
Heling & Shapira (1978)	Israel	R	17	T	118	C&R	S				χ^2
Jokinen et al. (1978)	Scandinavia	R	45	Ro	2459	C&R	S	✓			χ^2
Kerekes (1978)	Scandinavia	R	-	Ro	379	Ra	S	✓			χ^2
Kerekes (1978)	Scandinavia	R	-	Ro	188	Ra	S	✓			χ^2
Soltanoff (1978)	USA	R	-	T	266	Ra	L				χ^2
Heling & Kischinsky (1979)	Switzerland	R	13	Ro	202	C&R	L				χ^2
Barbakow et al. (1980a,b, 1981)	South Africa	R	60	T	335	C&R	L				χ^2
Cvek et al. (1982)	Scandinavia	R	83	Ro	45	Ra	S				χ^2
Nelson (1982)	UK	R	-	T	299	C&R	L				χ^2
Boggia (1983)	UK	R	-	T	52	Ra	S				χ^2
Klevant & Eggink (1983)	Holland	R	76	T	319	Ra	S				χ^2
Morse et al. (1983a,b,c)	USA	R	-	Ro	458	C&R	L				χ^2
Oliet (1983)	USA	C	-	T	338	C&R	L				χ^2
Swartz et al. (1983)	USA	R	-	Ro	1770	C&R	L				χ^2
Pekruhn (1986)	Saudi Arabia	R	81	T	925	C&R	S	✓			χ^2
Byström et al. (1987)	USA	C	56	Ro	79	Ra	S				χ^2
Halse & Molven (1987)	Scandinavia	R	63	Ro	551	Ra	S				χ^2
Matsumoto et al. (1987)	Japan	R	38	T	85	C&R	L		✓		χ^2
Ørstavik et al. (1987) & Eriksen et al. (1988)	Scandinavia	RCT	36	Ro	289	Ra	L		✓		RIDIT
Safavi et al. (1987)	USA	R	-	T	464	C&R	S	✓	✓		χ^2

Akerblom & Hasselgren (1988)	Scandinavia	R	73			Ro	64	C&R	S	✓			
Shah (1988)	India	C	70		✓	T	65	C&R	L	✓			LR
Sjögren et al. (1990)	Scandinavia	R	46			Ro	573	Ra	S		✓		
Murphy et al. (1991)	USA	R	-		✓	T	89	Ra	S				χ^2
Cvek (1992)	Scandinavia	R	76		✓	Ro	610	Ra	S				χ^2
Reid et al. (1992)	Australia	RCT	44		✓	Ro	74	C&R	S	✓			
Jurcak et al. (1993)	USA	R	58		✓	T	102	C&R	L	✓			
Smith et al. (1993)	UK	R	54			T	821	C&R	L				χ^2
Peak (1994)	UK	R	-			T	136	C&R	S		✓		χ^2
Friedman et al. (1995)	Canada	C	78			T	250	C&R	S				χ^2
Çalişken & Şen (1996)	Turkey	R	-		✓	T	172	C&R	S				χ^2
Østavik (1996)	Scandinavia	C	81			Ro	599	Ra	L	✓			RIDIT
Peretz et al. (1997)	Israel	R	-		✓	T	28	C&R	S	✓			χ^2
Sjögren et al. (1997)	Scandinavia	C	96			Ro	53	Ra	S	✓			
Lilly et al. (1998)	USA	R	-			T	22	C&R	S	✓			χ^2
Trope et al. (1999)	USA	RCT	-			T	102	Ra	S	✓			χ^2
Ricucci et al. (2000)	Italy	R	-			T	110	Ra	S	✓			χ^2
Weiger et al. (2000)	Germany	RCT	92			T	67	C&R	S	✓			LR
Chugal et al. (2001)	USA	R	75		✓	R	322	Ra	S	✓			LR
Deutsch et al. (2001)	USA	R	42			T	153	C&R	L	✓			χ^2
Heling et al. (2001)	Israel	R	-		✓	T	319	Ra	L	✓			ANOVA
Peak et al. (2001)	UK	R	-			T	406	C&R	L	✓			
Pettiette et al. (2001)	USA	RCT	66			T	40	Ra	L	✓			χ^2
Benenati & Khajotia (2002)	USA	R	29			T	894	Ra	S	✓			χ^2
Cheung (2002)	Hong Kong	R	28		✓	T	282	C&R	S	✓			Survival
Hoskinson et al. (2002)	UK	R	42		✓	Ro	413	C&R	S	✓			GEE
Peters & Wesselink (2002)	Holland	RCT	100			Ro	38	C&R	S	✓			χ^2

^aR, retrospective study; C, prospective cohort study; RCT, randomized controlled trial.

^bT, teeth; Ro, root (unit of measure was recorded as 'root' for those studies which has only included single rooted teeth in their sample).

^cC&R, combined clinical and radiographic examination; Ra, radiographic examination only.

^dS, strict criteria; L, loose criteria.

^eLR, single level logistic regression; GEE, generalized estimating equations; χ^2 , chi-squared test; RIDIT, relative incidence distribution; ANOVA, analysis of variance; Survival, survival analysis.

Table 3 Clinical prognostic factors included in studies

Author (year)	Gender	Age	Health	Tooth type	Pulpal status	Periapical status	Lesion size	Rubber dam	Obstruction	Apical size	Canal taper	Irrigant	Medicament	Culture test	RF material technique	Sealer	RF extent	Quality of RF	Acute flare up	Apical disturbance	Visits of treatment	Restoration	Abutment
Blayney (1922)					✓	✓		✓				✓									✓		✓
Auerbach (1938)					✓	✓		✓				✓									✓		
Buchbinder (1941)					✓	✓		✓				✓									✓		
Morse & Yates (1941)					✓	✓		✓				✓									✓		
Castagnola & Orlay (1952)					✓	✓		✓				✓									✓		
Grahnén & Hansen (1961)					✓	✓		✓				✓									✓		
Seltzer et al. (1963)		✓			✓	✓		✓				✓									✓		
Zeldow & Ingie (1963)					✓	✓		✓				✓									✓		
Bender et al. (1964)					✓	✓		✓				✓									✓		
Engström & Lundberg (1965)					✓	✓		✓				✓									✓		
Harty et al. (1970)		✓			✓	✓		✓				✓									✓		
Heling & Tamshe (1970, 1971)					✓	✓		✓				✓									✓		
Cvek (1972)					✓	✓	✓	✓				✓									✓		
Selden (1974)					✓	✓		✓				✓									✓		
Werts (1975)					✓	✓		✓				✓									✓		
Adenubi & Rule (1976)		✓			✓	✓		✓				✓									✓		
Heling & Shapira (1978)		✓			✓	✓		✓				✓									✓		
Jokinen et al. (1978)					✓	✓		✓				✓									✓		
Kerkes (1978)					✓	✓		✓				✓									✓		
Kerkes (1978)					✓	✓		✓				✓									✓		
Soltanoff (1978)					✓	✓		✓				✓									✓		
Heling & Kischinsky (1979)					✓	✓		✓				✓									✓		
Barbakow et al. (1980a,b, 1981)					✓	✓		✓	✓			✓									✓		
Cvek et al. (1982)					✓	✓		✓				✓									✓		
Nelson (1982)					✓	✓		✓				✓									✓		
Boggia (1983)					✓	✓		✓				✓									✓		
Klevant & Eggink (1983)					✓	✓		✓				✓									✓		
Morse et al. (1983a,b,c)					✓	✓		✓				✓									✓		
Oliet (1983)					✓	✓		✓				✓									✓		
Swartz et al. (1983)		✓			✓	✓		✓				✓									✓		
Pekruhn (1986)					✓	✓		✓				✓									✓		
Byström et al. (1987)					✓	✓		✓				✓									✓		
Halse & Molven (1987)					✓	✓		✓				✓									✓		
Matsumoto et al. (1987)					✓	✓		✓				✓									✓		

	8	13	4	13	4	6	33	2	1	2	32	20	14	38	25	29	7	0	5	35	8	2	
Ørstavik et al. (1987) &																							
Eriksen et al. (1988)																						✓	
Safavi et al. (1987)											✓											✓	
Akerblom &								✓														✓	
Hasselgren (1988)																						✓	
Shah (1988)																						✓	
Sjögren et al. (1990)																						✓	
Murphy et al. (1991)																						✓	
Cvek (1992)																						✓	
Reid et al. (1992)																						✓	
Jurcak et al. (1993)																						✓	
Smith et al. (1993)																						✓	
Peak (1994)																						✓	
Friedman et al. (1995)																						✓	
Çalışkan & Şen (1996)																						✓	
Ørstavik (1996)																						✓	
Peretz et al. (1997)																						✓	
Sjögren et al. (1997)																						✓	
Lilly et al. (1998)																						✓	
Trope et al. (1999)																						✓	
Ricucci et al. (2000)																						✓	
Weiger et al. (2000)																						✓	
Chugal et al. (2001)																						✓	
Deutsch et al. (2001)																						✓	
Heling et al. (2001)																						✓	
Peak et al. (2001)																						✓	
Pettiette et al. (2001)																						✓	
Benenati & Khajotia (2002)																						✓	
Cheung (2002)																						✓	
Hoskinson et al. (2002)																						✓	
Peters & Wesselink (2002)																						✓	
Total	8	13	4	13	4	6	33	2	1	2	32	20	14	38	25	29	7	0	5	35	8	2	

RF, root filling.

Table 4 Estimated success rates by study characteristics

Factor/categories	No. studies identified	Strict radiographic criteria				Loose radiographic criteria			
		No. studies	No. units	Estimated pooled success rates		No. studies	No. units	Estimated pooled success rates	
				^c Un-weighted (%)	^d Weighted (%)			^c Un-weighted (%)	^d Weighted (%)
Outcome measure used									
Radiographic	27	17	4745	74.4	74.1 (66.9–81.3)	14 (13) ^b	7177	83.4	84.1 (79.0–89.3)
Clinical + radiographic	36	23	10799	72.8	75.0 (68.4–81.7)	24 (23) ^b	10430	82.0	85.8 (81.8–89.9)
Duration after treatment (months)									
6	4	2	1120	17.3	29.6 (14.2–73.3)	2	4633	89.1	89.1 (78.8–99.5)
12	9	5	2080	68.6	67.7 (39.0–96.4)	4	798	76.7	69.5 (52.3–86.6)
24	6	3	2328	75.8	67.3 (43.8–90.9)	3	1103	82.7	82.5 (75.2–89.8)
36	5	2	941	80.4	80.6 (78.0–83.1)	3	254	69.3	66.6 (36.1–95.9)
48	6	5	2931	84.5	83.8 (79.3–88.3)	2	301	93.5	61.7 (25.0–96.0)
>48	9	8	1162	86.8	85.4 (80.3–90.6)	1	821	84.3	84.3 (81.8–86.8)
Year of publication									
Before 1960	5	2	1245	68.2	68.2 (65.6–70.8)	4	1580	82.2	84.2 (72.2–96.1)
1960s	5	2	1458	81.3	79.7 (74.0–85.4)	3	3669	80.4	80.4 (79.2–81.7)
1970s	12	9	5468	69.8	79.0 (66.7–91.3)	6 (5) ^b	4400	77.0	84.0 (71.3–96.6)
1980s	16	9	2834	77.8	74.8 (62.5–87.0)	10 (9) ^b	3770	89.7	88.2 (85.0–91.4)
1990s	14	10	2007	83.9	76.9 (69.7–84.1)	8	2271	83.3	85.5 (80.2–90.9)
2000s	11	8	2532	65.2	68.0 (60.5–75.4)	7	1917	85.2	85.1 (78.0–92.3)
Geographic location of study									
USA or Canada	24	9	2412	67.5	74.1 (64.9–83.2)	20 (19) ^b	9393	86.5	88.1 (84.9–91.2)
Scandinavian country	15	12	6435	71.5	80.5 (71.1–89.8)	3 (2) ^b	3347	70.4	70.3 (61.3–79.2)
Other country	24	19	6697	76.9	71.2 (64.4–78.1)	15	4867	83.4	84.5 (80.4–88.5)
Qualification of operators									
Undergraduate students	21	14	8306	68.4	74.8 (67.0–82.7)	11 (10) ^b	7808	79.9	83.3 (75.8–90.9)
GDP	7	6	1353	64.4	65.7 (56.3–75.1)	5	1228	85.5	86.2 (82.9–89.5)
Postgraduate students	4	4	1336	82.9	77.2 (64.5–89.8)	2	959	93.1	93.1 (91.5–94.7)
Specialist	23	11	3288	87.6	84.8 (80.1–89.4)	17 (16) ^b	6368	84.7	87.6 (83.9–91.3)

GDP, general dental practitioners.

^aTotal number of studies identified for the respective study characteristics is equal to or smaller than the summation of number of studies under strict and loose criteria as some studies reported success rates based on both criteria.

^bNumber in bracket indicating the number of studies included in the meta-analysis after those studies with 100% rates by the respective factor under investigation have been excluded.

^cUn-weighted pooled success rates were estimated based on the Hedges & Friedman (1997)'s approach.

^dWeighted pooled success rates were estimated using random effects meta-analysis (where there was only one study, its reported success rate and confident intervals were presented).




Figure 1 Probability of success based on strict radiographic criteria.




Figure 2 Probability of success based on loose radiographic criteria.

out in Scandinavian (15 studies, Sweden/Norway) or other countries (24 studies) including: UK (eight studies), Israel (four studies), Holland (two studies),

Switzerland (one study), Australia (one study), Germany (one study), Hong Kong (one study), India (one study), Italy (one study), Japan (one study), Saudi

Arabia (one study), South Africa (one study) and Turkey (one study) (Table 2). The studies performed in the North American countries reported the treatment outcome data more frequently based on loose radiographic criteria than on strict criteria. In contrast, most of the outcome data from the Scandinavian countries were based on strict rather than loose criteria. Based on the loose criteria, the pooled estimate of success rate of treatment carried out in Scandinavian countries (70.3%) was much lower than for those in North American (88.1%) or other (84.5%) countries; however, the pooled estimate for the Scandinavian countries, only consisted of two studies. In stark contrast, the pooled estimate of success rate from outcome data based on strict criteria from the Scandinavian countries (80.5%) was the highest (Table 4).

Qualification of operators (undergraduate, postgraduate, GDP and specialist)

Only two studies compared the outcome of root canal treatment by qualification of operators. Ingle *et al.* (1965) (a study excluded from this review; Table 1), found no significant difference in success rates of treatment carried out by undergraduates or private practitioners, in agreement with Cheung (2002) who reported the qualification and experience of operator had no influence on treatment outcome.

The majority of the reviewed studies classified operator qualification as: undergraduate students (21 studies), GDP (seven studies), postgraduate students (four studies) or specialists (23 studies). In five studies, treatment was carried out by a mixed group of operators and three studies did not provide this information. From the results, treatment carried out by postgraduate students and specialists had the highest weighted pooled estimate of success, regardless of strict or loose criteria (Table 4).

Source of heterogeneity

As the radiographic criteria for success have already been shown to have a significant effect on the pooled success rates, further meta-regression analyses were therefore carried out, separately on success rates based on strict or loose criteria, to explore which of the other study characteristics were potentially responsible for the statistical heterogeneity. None had significant effects on the success rates reported by the studies or could account for the heterogeneity (Table 5) in estimating the pooled success rate of primary root canal treatment.

Table 5 Results of meta-regression analysis to account for the source of heterogeneity

Covariate included	Strict		Loose	
	I^2	τ^2	I^2	τ^2
No. covariate included	0.985	0.0247	0.973	0.0085
Year of publication (before 1970s, 1970–1989, 1990–2002)	0.983	0.0265	0.971	0.0098
Geographical location of study (USA, Scandinavian or other countries)	0.984	0.0244	0.952	0.0069
Unit of measure (root or tooth)	0.984	0.0253	0.961	0.0081
Qualification of operator (specialist, postgraduate, undergraduate, GDP or mixed group)	0.979	0.0209	0.974	0.0073
Criteria for success (radiographic vs. combined radiographic & clinical)	0.986	0.0254	0.974	0.0088
Duration after treatment (at least 4 years or shorter)	0.985	0.0228	0.973	0.0085
Recall rate	0.986	0.0218	0.975	0.0117

GDP, general dental practitioners; I^2 , proportion of total variation due to heterogeneity across studies; τ^2 , estimate of between-study variance (if the I^2 and τ^2 values were reduced by 10% after including a covariate in the regression model as compared with the values estimated without any covariates entered, the respective covariate was considered to be a potential source of heterogeneity).

Discussion

Most of the selected studies were prospective cohort or retrospective studies, therefore the levels of evidence provided by them are grade B (levels 2 or 3) based on the criteria given by the Oxford centre for evidence-based medicine (Phillips *et al.* 1998 <http://www.cebm.net/index.aspx?0=1025>). There were only six randomized trials investigating different aspects of root canal treatment procedures on outcomes, including the effects of sealers (AH26 [DeTrey AG, Zurich, Switzerland]; Procosol [Star Dental, Conshohocken, PA, USA], Kloropercha) (Ørstavik *et al.* 1987); root filling materials (Hydron[®] [Hydron Technologies, Pompano Beach, FL, USA], gutta-percha with AH26[®] sealer) (Reid *et al.* 1992); single-visit versus multiple-visit root canal treatment (Trope *et al.* 1999, Weiger *et al.* 2000, Peters & Wesselink 2002); use of Ca(OH)₂ dressing versus no medicament in multiple-visit treatment (Trope *et al.* 1999); and the use of stainless steel versus nickel–titanium hand files (Pettiette *et al.* 2001).

The Cochrane Oral Health group's current guidelines for a systematic review states that 'the scope of the review is to include all RCTs, where RCTs are inappropriate, rather than unavailable, other levels of evidence may be considered' (http://www.ohg.cochrane.org/forms/writing_review.pdf September 2006). However, the authors decided that the numerous observational studies, whilst not having the feature of randomization or control groups, represented useful and useable data that could not be deemed inferior by any other criteria. Instead of using exclusion rules to control the heterogeneity of design, this systematic review followed the recommendation by Stroup *et al.* (2000). Broad inclusion criteria for studies were used and analyses were performed to investigate the effect of study characteristics on the estimated pooled success rates. Despite using broad inclusion criteria, several well designed and executed studies such as those by Strindberg (1956), Ørstavik and Hörsted-Bindslev (1993); Ørstavik *et al.* (1986) and others (Table 1) had to be excluded.

The goal was to explore the available data and partition it to reveal the effect of study characteristics, general patient factors, individual pre-, intra-, and postoperative factors on treatment outcome, whilst triangulating the outcomes of different approaches of exploration.

Preliminary data collection was carried out by two authors (Y-LN and KG) to explore its diversity. Amongst the studies reviewed, there were substantial variations in study characteristics such as sample selection, definition of successful cases, duration after treatment, type and strategy of data collection as well as data analyses. Some of the potential clinical prognostic factors (tooth type, age grouping, size of apical preparation, definition of apical disturbance, apical extent of root filling, quality of root filling and quality of coronal restoration) were sub-classified differently between studies. Therefore, a strategy for pooling the data by recalculation of the available figures was derived. Based on this strategy, a data collection form was designed, tested, refined and adopted. Despite this, disagreement amongst the reviewers existed ($\kappa = 0.57-0.61$). The disagreement could be traced to a lack of clarity in the presentation of methodology and results. For some studies, data had to be extracted from the discussion section where it was sometimes first introduced. When there was a disagreement, an agreement was negotiated between the examiners by presenting the case for each view. In the majority of

cases, the source of errors were easily identified and corrected.

The un-weighted pooled success rate by each factor was calculated based on the approach used by Hepworth & Friedman (1997). However, this approach does not take into consideration, the within and between study variations as opposed to the study-design-specific weighted pooled success rates estimated using random effects meta-analysis. The discrepancies in the success rates estimated using the two approaches are well demonstrated in Table 4. Therefore, the results based on the un-weighted pooled success rates were not considered in the following discussions. The estimation of pooled success rates for some sub-group analyses within some study characteristics were based on small data sets, restricting their value.

The significant difference in success rates judged by strict or loose radiographic criteria has already been iterated but in addition, the data based on strict criteria revealed a clear trend for differences in the pooled success rates from studies adopting different follow-up durations; for example, 6 month follow-up compared with longer duration. It should be mandatory to state the criteria for success as part of the methodology of future clinical root canal treatment outcome studies, preferably stratified by both loose and strict radiographic criteria. The European Society of Endodontology's (2006) suggest a clinical and radiographic follow-up after at least 1 year with annual recall for up to 4 years before a case is judged a failure. The American Association of Endodontists suggests clinical and radiographic evaluation for a 4- to 5-year period, with the additional proviso of determining the functionality of the treated tooth (<http://www.aae.org/dentalpro/guidelines.htm>). The origin of this is probably based on the work of Strindberg (1956). From a research perspective and based on this review, the cases should be reviewed for a minimum of 1 year and preferably for at least 3 years, after completion of treatment. It would be preferable to standardize the duration after treatment for all the patients or at least to include the duration as a covariate into the statistical model to account for any variations in the success rate because of the different follow-up times. The best choice of statistical analysis would be to analyse time to healing (success) with survival analysis techniques, however, it would require regular follow-up of all patients. The reality is that the longer the duration of follow-up after treatment, the greater the drop-out rate at recall. Therefore, a balance has to be struck between these competing ideals in both the medical and dental

fields, although the use of financial incentives may help improve recall rates (Wang *et al.* 2004).

This review also highlighted two important methodological shortcomings in published root canal treatment outcome studies. The variability in radiographic assessment because of subjectivity in radiograph reading is well recognized (Goldman *et al.* 1972), yet the good practice of employing at least two pre-calibrated observers with intra- & inter-observer agreement tests, was not adopted by most of the studies ($n = 56$). In addition, the statistical methods used for analysing the association between potential influencing factors and treatment outcome, did not take account of the effects of potential confounders.

The overall success rates were not affected by 'year of publication' or 'geographic location of study'. In the former category, the measure of relevance should really be the 'year in which treatment was carried out' but few studies provide this information. Nevertheless, the absence of obvious improvement in success rates by the year of publication suggests that the advances in technology and materials used for root canal treatment do not appear to have influenced treatment outcome significantly. Such a suggestion is strongly refuted by endodontists on the grounds that the apparent lack of improvement in success rates is a function of more adventurous case selection fuelled by confidence in better skills and outcomes. The validity of this proposition is explored further in the second part. For the present, it is argued that the lack of improvement in success rates could be attributed to the fact that, whilst technology has improved instruments and materials to achieve a set of goals, the principles underpinning those goals have not changed over the duration covered by this review (Hall 1928). This brings to the fore, the classic debate about the relative value of biological versus the technical principles in dentistry (Noyes 1922, Naidorf 1972). Noyes (1922) lamented that dentists were not trained to think in biological concepts but to act in mechanical procedures; whilst Naidorf (1972) applauding the technical excellence achieved by the pre-occupation of dentists with this element, deplored the lack of biological awareness of the basic pathology of the problem or the biological consequences of the treatment. The clinical academics in this discipline would probably sustain the validity of these assertions, even today. It is interesting to note that the success rates of studies from the North American countries, where the use of contemporary technology is probably most widely recommended fared no better than those from other countries. Further-

more, the adoption of strict radiographic criteria and microbiological awareness in their approach appeared to bring better results in studies performed in the Scandinavian countries. This speculation is important because it centres around the debates that raged in the 1960s and 1970s about the value of the microbial culture test in informing the progress of treatment, a practice, long as abandoned as unnecessary (Engström *et al.* 1964, Mikkelsen & Theilade 1969, Oliet & Sorin 1969, Morse 1971, Sims 1973, Frank *et al.* 1978, Molander *et al.* 1996a,b). This ultimately led to the adoption of single-visit treatment by many endodontists on the basis of the cost-benefit analysis (Spångberg 2001) an issue that will be explored further in the second part. The historical importance of this biological versus technical debate is important to appreciate, because it fundamentally changed the way root canal treatment was conceived and practiced; from a microbially aware post-focal infection era, to one dominated by a technological awareness but relative microbiological ignorance. The problem of geographical location also merits close inspection, as sometimes, a single study may report pooled data from multi-centre evaluations (Friedman *et al.* 1995).

Although the educational and experience background of the operators had no significant influence on their respective success rates in individual studies, the estimated pooled success rates for endodontists or postgraduates were higher than for other dentist groups in this review. The important influence of the background of operators on the technical outcome of endodontic procedures has been demonstrated in laboratory studies (Gulabivala *et al.* 2000, Van Zyl *et al.* 2005) but there is a lack of appropriate tools or methodology to objectively quantify operator skills. The role of such refined technical skills must surely be balanced against the overall understanding of the problem and the motivation and integrity with which the procedure is performed.

Conclusion

The estimated weighted pooled success rates of treatments completed at least 1 year previously, ranged between 68% and 85% when strict criteria were used. The reported success rates have failed to improve over the last four or five decades. The quality of evidence for treatment factors affecting primary root canal treatment outcome is sub-optimal; there was substantial variation in the study-designs. It would be desirable to standardize aspects of study-design, data recording and

presentation format of outcome data in the much needed future outcome studies. The second part of this paper will present the results of meta-analyses and meta-regression to investigate the effect of individual clinical factors on the success rates of primary root canal treatment.

Acknowledgement

The authors acknowledge the mentorship provided by Professor Mark Gilthorpe of the University of Leeds (previously based at the Eastman Dental Institute) during the early stages of this project.

References

- Adenubi JO (1978) The criteria for success in endodontics. *Nigerian Medical Journal* **8**, 404–10.
- Adenubi JO, Rule DC (1976) Success rate for root fillings in young patients. *British Dental Journal* **141**, 237–41.
- Akerblom A, Hasselgren G (1988) The prognosis for endodontic treatment of obliterated root canals. *Journal of Endodontics* **14**, 565–7.
- Appleton JLT (1932) A note on the clinical value of bacteriological controlling the treatment of periapical infection. *Dental Cosmos* **IXXiv**, 798–800.
- Ashkenaz PJ (1984) One-visit endodontics. *Dental Clinics North America* **28**, 853–63.
- Auerbach MB (1938) Clinical approach to the problem of pulp canal therapy. *Journal of American Dental Association* **25**, 939–42.
- Augsburger RA, Peters DD (1990) Radiographic evaluation of extruded obturation materials. *Journal of Endodontics* **16**, 492–7.
- Barbakow FH, Cleaton-Jones P, Friedman D (1980a) An evaluation of 566 cases of root canal therapy in general dental practice. 1. Diagnostic criteria and treatment details. *Journal of Endodontics* **6**, 456–60.
- Barbakow FH, Cleaton-Jones P, Friedman D (1980b) An evaluation of 566 cases of root canal therapy in general dental practice. 2. Postoperative observations. *Journal of Endodontics* **6**, 485–9.
- Barbakow FH, Cleaton-Jones PE, Friedman D (1981) Endodontic treatment of teeth with periapical radiolucent areas in a general dental practice. *Oral Surgery, Oral Medicine, Oral Pathology* **51**, 552–9.
- Basmadjian-Charles CL, Farge P, Bourgeois DM, Lebrun T (2002) Factors influencing the long-term results of endodontic treatment: a review of the literature. *International Dental Journal* **52**, 81–6.
- Bender IB, Seltzer S, Turkenkopf S (1964) To culture or not to culture? *Oral Surgery, Oral Medicine, Oral Pathology* **18**, 527–40.
- Benenati FW, Khajotia SS (2002) A radiographic recall evaluation of 894 endodontic cases treated in a dental school setting. *Journal of Endodontics* **28**, 391–5.
- Blayney JR (1922) The clinical results of pulp treatment. *Journal of National Dental Association* **16**, 198–208.
- Boggia R (1983) A single-visit treatment of septic root canals using periapically extruded endomethasone. *British Dental Journal* **155**, 300–5.
- Buchbinder M (1936) A statistical study of root-canal therapy. *The Dental Cosmos* **78**, 20–6.
- Buchbinder M (1941) A statistical comparison of cultured and non-cultured root canal cases. *Journal of Dental Research* **2**, 93–6.
- Byström A, Happonen RP, Sjögren U, Sundqvist G (1987) Healing of periapical lesion of pulpless teeth after endodontic treatment with controlled asepsis. *Endodontics & Dental Traumatology* **3**, 58–63.
- Çalışken MK, Şen BH (1996) Endodontic treatment of teeth with apical periodontitis using calcium hydroxide: a long-term study. *Endodontics & Dental Traumatology* **12**, 215–21.
- Caplan DJ, White BA (2001) Clinical factors related to noncompletion of root canal therapy. *Journal of Public Health in Dentistry* **61**, 6–13.
- Caplan DJ, Kolker J, Rivera EM, Walton RE (2002) Relationship between number of proximal contacts and survival of root canal treated teeth. *International Endodontic Journal* **35**, 193–9.
- Castagnola L, Orlay HG (1952) Treatment of gangrene of the pulp by the Walkhoff method. *British Dental Journal* **19**, 93–100.
- Cheung GS (2002) Survival of first-time nonsurgical root canal treatment performed in a dental teaching hospital. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **93**, 596–604.
- Chugal NM, Clive JM, Spångberg LSW (2001) A prognostic model for assessment of the outcome of endodontic treatment: effect of biologic and diagnostic variables. *Oral Surgery, Oral Medicine, Oral Pathology, Radiology and Endodontics* **91**, 342–52.
- Cochran WG (1954) The combination of estimates from different experiments. *Biometrics* **10**, 101–29.
- Coolidge ED (1926) A discussion of clinical results of root-canal treatment and filling. *The Dental Cosmos* **IXix**, 1280–8.
- Curson I (1966) Endodontic techniques – prognosis in endodontics. *British Dental Journal* **20**, 568–70.
- Cvek M (1972) Treatment of non-vital permanent incisors with calcium hydroxide. *Odontologisk Revy* **23**, 27–44.
- Cvek M (1992) Prognosis of luxated non-vital maxillary incisors treated with calcium hydroxide and filled with gutta-percha. A retrospective clinical study. *Endodontics & Dental Traumatology* **8**, 45–55.
- Cvek M, Hollender L, Nord C-E (1976) Treatment of non-vital permanent incisors with calcium hydroxide. *Odontologisk Revy* **27**, 93–108.

- Cvek M, Granath L, Lundberg M (1982) Failures and healing in endodontically treated non-vital anterior teeth with posttraumatically reduced pulpal lumen. *Acta Odontologica Scandinavica* **40**, 223–8.
- DerSimonian R, Laird N (1986) Meta-analysis in clinical trials. *Control Clinical Trials* **7**, 177–88.
- Deutsch AS, Musikant BL, Cohen BI, Kase D (2001) A study of one visit treatment using EZ-fill root canal sealer. *Endodontic Practice* **4**, 29–36.
- Engström B, Lundberg M (1965) The correlation between positive culture and the prognosis of root canal therapy after pulpectomy. *Odontologisk Revy* **16**, 193–204.
- Engström B, Segerstad LHA, Ramstrom G, Frostell G (1964) Correlation of positive cultures with the prognosis for root canal treatment. *Odontologisk Revy* **15**, 257–70.
- Eriksen HM, Ørstavik D, Kerekes K (1988) Healing of apical periodontitis after endodontic treatment using three different root canal sealers. *Endodontics & Dental Traumatology* **4**, 114–7.
- European Society of Endodontology (2006) Quality guidelines for endodontic treatment: consensus report of the European Society of Endodontology. *International Endodontic Journal* **39**, 921–30.
- Eysenck HJ (1994) Meta-analysis and its problems. *British Medical Journal* **309**, 789–92.
- Frank AL, Abou Rass M, Glick DH (1978) Changing trends in endodontics. *Journal of American Dental Association* **96**, 202–10.
- Friedman S (1997) Success and failure of initial endodontic therapy. *Ontario Dentist* **74**, 35–8.
- Friedman S, Löst C, Zarrabian M, Trope M (1995) Evaluation of success and failure after endodontic therapy using glass ionomer cement sealer. *Journal of Endodontics* **21**, 384–90.
- Frostell G (1963) Clinical significance of the root canal culture. *Transactions of Third International Conference on Endodontics*, 112–22.
- Goldman M, Pearson AH, Darzenta N (1972) Endodontic success – who's reading the radiograph?. *Oral Surgery, Oral Medicine, Oral Pathology* **33**, 432–7.
- Grahnen H, Hansen L (1961) The prognosis of pulp and root canal therapy. *Odontologisk Revy* **12**, 146–65.
- Grossman LI, Shepard LI, Person LA (1964) Roentgenologic and clinical evaluation of endodontically treated teeth. *Oral Surgery* **17**, 368–74.
- Grove CJ (1921) Nature's method of making perfect root fillings following pulp removal, with a brief consideration of the development of secondary cementum. *The Dental Cosmo* **11**, 698–981.
- Grove CJ (1923) A method of pulp removal to prevent periapical infection. *Journal of American Dental Association* **10**, 93–108.
- Gulabivala K, Abdo S, Sherriff M, Regan JD (2000) The influence of interfacial forces and duration of filing on root canal shaping. *Endodontics & Dental Traumatology* **16**, 166–74.
- Gutknecht N, Kaiser F, Hassan A, Lampert F (1996) Long-term clinical evaluation of endodontically treated teeth by Nd:YAG lasers. *Journal of Clinical Laser Medicine and Surgery* **14**, 7–11.
- Hall EM (1928) Pulpless teeth. *The Dental Cosmo* **Ixx**, 145–50.
- Halse A, Molven O (1987) Overextended gutta-percha and Kloroperka N-Ø root canal fillings. Radiographic findings after 10–17 years. *Acta Odontologica Scandinavica* **45**, 171–7.
- Harty FJ, Parkins BJ, Wengraf AM (1970) Success rate in root canal therapy. A retrospective study on conventional cases. *British Dental Journal* **128**, 65–70.
- Heling B, Kischinovsky D (1979) Factors affecting successful endodontic therapy. *Journal of British Endodontic Society* **12**, 83–9.
- Heling B, Shapira J (1978) Roentgenologic and clinical evaluation of endodontically treated teeth, with or without negative culture. *Quintessence International* **11**, 79–84.
- Heling B, Tamshe A (1970) Evaluation of the success of endodontically treated teeth. *Oral Surgery, Oral Medicine, Oral Pathology* **30**, 533–6.
- Heling B, Tamshe A (1971) Statistical evaluation of the success of endodontically treated teeth. *New York Journal of Dentistry* **41**, 69–82.
- Heling I, Bialla-Shenkman S, Turetzky A, Horwitz J, Sela J (2001) The outcome of teeth with periapical periodontitis treated with nonsurgical endodontic treatment: a computerized morphometric study. *Quintessence International* **32**, 397–400.
- Hepworth MJ, Friedman S (1997) Treatment outcome of surgical and non-surgical management of endodontic failures. *Journal of Canadian Dental Association* **63**, 364–71.
- Hession RW (1981) Long-term evaluation of endodontic treatment: anatomy, instrumentation, obturation – the endodontic practice triad. *International Endodontic Journal* **14**, 179–84.
- Hinman TP (1921) The interpretation of X-ray pictures of apical granulations, giving differential diagnosis of cases favourable and cases unfavourable for treatment and root-canal filling. *Journal of National Dental Association* **8**, 83–7.
- Hoskinson SE, Ng YL, Hoskinson AE, Moles DR, Gulabivala K (2002) A retrospective comparison of outcome of root canal treatment using two different protocols. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **93**, 705–15.
- Ingle JL, Beveridge EE, Glick DH, Weichman JA (1965) Modern endodontic therapy. In: Ingle JL, Bakland LK, eds. *Endodontics*, 4th edn. Baltimore, MA: Williams and Wilkins 1994, pp. 27–53.
- Jokinen MA, Kotilainen R, Poikkeus P, Poikkeus R, Sarkki L (1978) Clinical and radiographic study of pulpectomy and root canal therapy. *Scandinavian Journal of Dental Research* **86**, 366–73.

- Jurcak JJ, Bellizzi R, Loushine RJ (1993) Successful single-visit endodontics during operation desert shield. *Journal of Endodontics* **19**, 412–3.
- Kerekes K (1978) Radiographic assessment of an endodontic treatment method. *Journal of Endodontics* **4**, 210–3.
- Kerekes K, Tronstad L (1979) Long-term results of endodontic treatment performed with a standardized technique. *Journal of Endodontics* **5**, 83–90.
- Klevant FJH, Eggink CO (1983) The effect of canal preparation on periapical disease. *International Endodontic Journal* **16**, 68–75.
- Kojima K, Inamoto K, Nagamatsu K et al. (2004) Success rate of endodontic treatment of teeth with vital and nonvital pulps. A meta-analysis. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **97**, 95–9.
- Kullendorff B, Grondahl K, Rohlin M, Henrikson CO (1988) Subtraction radiography for the diagnosis of periapical bone lesions. *Endodontics and Dental Traumatology* **4**, 253–9.
- Lazarski MP, Walker WA III, Flores CM, Schindler WG, Hargreaves KM (2001) Epidemiological evaluation of the outcomes of non-surgical root canal treatment in a large cohort of insured dental patients. *Journal of Endodontics* **27**, 791–6.
- Lewsey JD, Gilthorpe MS, Gulabivala K (2001) An introduction to meta-analysis within the framework of multilevel modeling using the probability of success of root canal treatment as an illustration. *Community Dental Health* **18**, 131–7.
- Lilly JP, Cox D, Arcuri M, Krell KV (1998) An evaluation of root canal treatment in patients who have received irradiation to the mandible and maxilla. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **86**, 224–6.
- Lynch CD, Burke FM, NíRiordáin R, Hannigan A (2002) Preoperative periapical status and survival of root filled teeth: a pilot investigation. *International Endodontic Journal* **35**, 492 (Abstract).
- Macphee GG (1936) The problem of the pulpless tooth. *British Dental Journal* **58**, 119–25.
- Markitziu A, Heling I (1981) Endodontic treatment of patients who have undergone irradiation of the head and neck. A longitudinal follow-up of eleven endodontically treated teeth. *Oral Surgery, Oral Medicine, Oral Pathology* **52**, 294–8.
- Matsumoto T, Nagai T, Ida Km Ito M et al. (1987) Factors affecting successful prognosis of root canal treatment. *Journal of Endodontics* **13**, 239–42.
- Mikkelsen L, Theilade E (1969) False negative cultures from root canals. Inhibition of bacterial growth by some drugs used in root canal therapy. *Acta Odontologica Scandinavica* **27**, 387–96.
- Molander A, Reit C, Dahlén G (1996a) Microbiological root canal sampling; diffusion of a technology. *International Endodontic Journal* **29**, 163–7.
- Molander A, Reit C, Dahlén G (1996b) Reasons for dentists' acceptance or rejection of microbiological root canal sampling. *International Endodontic Journal* **29**, 168–72.
- Molven O, Halse A (1988) Success rates for gutta-percha and Kloropercha N-Ø root fillings made by undergraduate students: radiographic findings after 10–17 years. *International Endodontic Journal* **21**, 243–50.
- Morse DR (1971) The endodontic culture technique: an impractical and unnecessary procedure. *Dental clinics of North America* **15**, 793–806.
- Morse FW, Yates MF (1941) Follow up studies of root-filled teeth in relation to bacteriologic findings. *Journal of American Dental Association* **28**, 956–71.
- Morse DR, Esposito JV, Pike C, Furst ML (1983a) A radiographic evaluation of the periapical status of teeth treated by the gutta-percha-eucapercha endodontic method: a one-year follow-up study of 458 root canals. *Part I. Oral Surgery, Oral Medicine, Oral Pathology* **55**, 607–10.
- Morse DR, Esposito JV, Pike C, Furst ML (1983b) A radiographic evaluation of the periapical status of teeth treated by the gutta-percha-eucapercha endodontic method: a one-year follow-up study of 458 root canals. *Part II. Oral Surgery, Oral Medicine, Oral Pathology* **56**, 89–96.
- Morse DR, Esposito JV, Pike C, Furst ML (1983c) A radiographic evaluation of the periapical status of teeth treated by the gutta-percha-eucapercha endodontic method: a one-year follow-up study of 458 root canals. *Part III. Oral Surgery, Oral Medicine, Oral Pathology* **56**, 190–7.
- Murakami M, Inoue S, Inoue N (2002) Clinical evaluation of audiometric control root canal treatment: a retrospective case study. *Quintessence International* **33**, 465–74.
- Murphy WK, Kaugars GE, Collett WK, Dodds RN (1991) Healing of periapical radiolucencies after nonsurgical endodontic therapy. *Oral Surgery, Oral Medicine, Oral Pathology* **71**, 620–4.
- Naidorf IJ (1972) Inflammation and infection of pulp and periapical tissues. *Oral Surgery, Oral Medicine, Oral Pathology* **34**, 486–97.
- Neiderman R, Theodosopoulou JN (2003) A systematic review of *in vivo* retrograde obturation materials. *International Endodontic Journal* **36**, 577–85.
- Nelson IA (1982) Endodontics in general practice – a retrospective study. *International Endodontic Journal* **15**, 168–72.
- Nichols E (1963) Assessment of the periapical status of pulpless teeth. *British Dental Journal* **114**, 453–9.
- Noyes FE (1922) (In Blayney JR). The clinical results of pulp treatment. *Journal of National Dental Association* **16**, 198–208.
- Oliet S (1983) Single-visit endodontics: a clinical study. *Journal of Endodontics* **9**, 147–52.
- Oliet S, Sorin SM (1969) Evaluation of clinical results based upon culturing root canals. *Journal of British Endodontic Society* **3**, 3–6.
- Oliver CM, Abbott PV (2001) Correlation between clinical success and apical dye penetration. *International Endodontic Journal* **34**, 637–44.
- Ørstavik D, Hörsted-Bindslev P (1993) A comparison of endodontic treatment results at two dental schools. *International Endodontic Journal* **26**, 348–54.

- Ørstavik D, Kerekes K, Eriksen HM (1986) The periapical index: a scoring system for radiographic assessment of apical periodontitis. *Endodontics & Dental Traumatology* **2**, 20–34.
- Ørstavik D, Kerekes K, Eriksen HM (1987) Clinical performance of three endodontic sealers. *Endodontics & Dental Traumatology* **3**, 178–86.
- Østavik D (1996) Time-course and risk analyses of the development and healing of chronic apical periodontitis in man. *International Endodontic Journal* **29**, 150–5.
- Paik S, Sechrist C, Torabinejad M (2004) Levels of evidence for the outcome of endodontic retreatment. *Journal of Endodontics* **30**, 745–50.
- Peak JD (1994) The success of endodontic treatment in general dental practice: a retrospective clinical and radiographic study. *Primary Dental Care* **1**, 9–13.
- Peak JD, Hayes SJ, Bryant ST, Dummer PM (2001) The outcome of root canal treatment. A retrospective study within the armed forces (Royal Air Force). *British Dental Journal* **190**, 140–4.
- Pekruhn RB (1986) The incidence of failure following single-visit endodontic therapy. *Journal of Endodontics* **12**, 68–72.
- Peretz B, Yakir O, Fuks AB (1997) Follow up after root canal treatment of young permanent molars. *Journal of Clinical Pediatric Dentistry* **21**, 237–40.
- Peters LB, Wesselink PR (2002) Periapical healing of endodontically treated teeth in one and two visits obturated in the presence or absence of detectable microorganisms. *International Endodontic Journal* **35**, 660–7.
- Peterson J, Gutmann JL (2001) The outcome of endodontic re-surgery: a systematic review. *International Endodontic Journal* **34**, 169–75.
- Pettiette MT, Delano O, Trope M (2001) Evaluation of success rate of endodontic treatment performed by students with stainless-steel K-files and nickel–titanium hand files. *Journal of Endodontics* **27**, 124–7.
- Phillips B, Ball C, Sackett D et al. (1998) *Oxford Centre for Evidence-Based Medicine Levels of Evidence in Level of Evidence and Grades of Recommendation*. Oxford centre for evidence-based medicine http://www.cebm.net/levels_of_evidence.asp#levels.
- Puterbaugh PG (1926) Pulp canal therapeutics. *Journal of American Dental Association* **Oct**, 1384–91.
- Ratliff DP (1973) The success rate of root canal filling in permanent incisors in children. *University of Newcastle Upon Tyne Medical Gazette* **67**, 42–5.
- Reid RJ, Abbott PV, McNamara JR, Heithersay GS (1992) A five-year study of hydron root canal fillings. *International Endodontic Journal* **25**, 213–20.
- Rhein ML, Krasnow F, Gies WJ (1926) A prolonged study of the electrolytic treatment of dental focal infection – a preliminary report. *The Dental Cosmo* **Ixviii**, 971–81.
- Ricucci D, Langeland K (1997) Incomplete calcium hydroxide removal from the root canal: a case report. *International Endodontic Journal* **30**, 418–21.
- Ricucci D, Grondahl K, Bergenholtz G (2000) Periapical status of root-filled teeth exposed to the oral environment by loss of restoration or caries. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **90**, 354–9.
- Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS (1996) Evidence based medicine: what it is and what it isn't. *British Medical Journal* **13**, 71–2.
- Safavi KE, Dowden WE, Langeland K (1987) Influence of delayed coronal permanent restoration on endodontic prognosis. *Endodontics & Dental Traumatology* **3**, 187–91.
- Sathorn C, Parashos P, Messer HH (2005) Effectiveness of single- versus multiple-visit endodontic treatment of teeth with apical periodontitis: a systematic review and meta-analysis. *International Endodontic Journal* **38**, 347–55.
- Selden HS (1974) Pulpoperiapical disease: diagnosis and healing. A clinical endodontic study. *Oral Surgery, Oral Medicine, Oral Pathology* **27**, 271–83.
- Seltzer S, Bender IB, Turkenkopf S (1963) Factors affecting successful repair after root canal therapy. *Journal of American Dental Association* **57**, 651–62.
- Seto BG, Beumer J, Kagawa T, Klokkevold P, Wolinsky L (1985) Analysis of endodontic therapy in patients irradiated for head and neck cancer. *Oral Surgery, Oral Medicine, Oral Pathology* **60**, 540–5.
- Shah N (1988) Non-surgical management of periapical lesions: a prospective study. *Oral Surgery, Oral Medicine, Oral Pathology* **66**, 365–71.
- Shi Y, Wang J, Cao C (1997) Clinical studies on pulpitis and periapical periodontitis caused by traumatic occlusion. *Chinese Journal of Stomatology* **32**, 23–5.
- Sims W (1973) Some comments on the microbiological examination of root canals. *Journal of Dentistry* **2**, 2–6.
- Sjögren U, Hägglund B, Sundqvist G, Wing K (1990) Factors affecting the long-term results of endodontic treatment. *Journal of Endodontics* **16**, 498–504.
- Sjögren U, Figdor D, Persson S, Sundqvist G (1997) Influence of infection at the time of root filling on the outcome of endodontic treatment of teeth with apical periodontitis. *International Endodontic Journal* **30**, 297–306.
- Smith CS, Setchell DJ, Harty FJ (1993) Factors influencing the success of conventional root canal therapy – a five-year retrospective study. *International Endodontic Journal* **26**, 321–33.
- Soltanoff W (1978) A comparative study of the single-visit and the multiple-visit endodontic procedure. *Journal of Endodontics* **4**, 278–81.
- Spångberg LS (2001) Evidence-based endodontics: the one-visit treatment idea. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **91**, 617–8.
- Stabholz A (1990) Success rate in endodontics. *Alpha Omegan* **83**, 20–4.
- Storms JL (1969) Factors that influence the success of endodontic treatment. *Journal of Canadian Dental Association* **35**, 83–97.

- Strindberg LZ (1956) The dependence of the results of pulp therapy on certain factors – an analytical study based on radiographic and clinical follow-up examinations. *Acta Odontologica Scandinavica* **14**, 1–175.
- Stroup DF, Berlin JA, Morton SC *et al.* (2000) Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *Journal of American Medical Association* **19**, 2008–12.
- Swartz DB, Skidmore AE, Griffin JA (1983) Twenty years of endodontic success and failure. *Journal of Endodontics* **9**, 198–202.
- Taintor JF, Biesterfeld RC, Bonness BW (1978) Evaluating clinical endodontic success. *Dental Practice* **Dec**, 36–41.
- Teo CS, Chan NC, Lim SS (1986) Success rate in endodontic therapy – a retrospective study. *Part I. Dental Journal of Malaysia* **9**, 7–10.
- Thoden van Velzen SK, Duivenvoorden HJ, Schuurs AH (1981) Probabilities of success and failure in endodontic treatment: a Bayesian approach. *Oral Surgery, Oral Medicine, Oral Pathology* **52**, 85–90.
- Thompson SG, Higgins JP (2002) How should meta-regression analyses be undertaken and interpreted? *Statistics in Medicine* **15**, 1559–73.
- Trope M, Delano O, Ørstavik D (1999) Endodontic treatment of teeth with apical periodontitis: single vs multivisit treatment. *Journal of Endodontics* **25**, 345–50.
- Van Zyl S, Gulabivala K, Ng Y-L (2005) The effect of customization of master gutta-percha on the apical control of root canal fillings. *International Endodontic Journal* **38**, 658–66.
- Vernieks AA, Messer LB (1978) Calcium hydroxide induced healing of periapical lesions: a study of 78 non-vital teeth. *Journal of British Endodontic Society* **11**, 61–8.
- Waltimo TM, Boiesen J, Eriksen HM, Ørstavik D (2001) Clinical performance of 3 endodontic sealers. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **92**, 89–92.
- Wang N, Knight K, Dao T, Friedman S (2004) Treatment outcome in endodontics – The Toronto Study. Phases I and II: apical surgery. *Journal of Endodontics* **30**, 751–61.
- Weiger R, Axmann-Krcmar D, Löst C (1998) Prognosis of conventional root canal treatment reconsidered. *Endodontics & Dental Traumatology* **14**, 1–9.
- Weiger R, Rosendahl R, Löst C (2000) Influence of calcium hydroxide intracanal dressings on the prognosis of teeth with endodontically induced periapical lesions. *International Endodontic Journal* **33**, 219–26.
- Weine FS, Buchanan SL (1997) Controversies in clinical endodontics: Part 2. Single-appointment vs multiple-appointment treatment. *The Compendium of Continuing Education in Dentistry* **18**, 140–4, 146, 148 *passim*; quiz 156.
- Werts R (1975) Endodontic treatment: a five-year follow-up. *Dental Practice* **Jan**, 29–30.
- Wong M, Shelley JJ, Bodey T, Hall R (1992) Delayed root canal therapy: an analysis of treatment over time. *Journal of Endodontics* **18**, 387–90.
- Zeldow BJ, Ingle JI (1963) Correlation of the positive culture to the prognosis of endodontically treated teeth: a clinical study. *Journal of American Dental Association* **66**, 23–7.